

On finite size corrections to the dispersion relations of giant magnon and single spike on γ -deformed $T^{1,1}$

M. Michalcik^{**} and R. C. Rashkov^{*, $\diamond$ \dagger}

^{*} Institute for Theoretical Physics,
Vienna University of Technology,
Wiedner Hauptstr. 8-10, 1040 Vienna, Austria,

$\diamond$ Department of Physics, Sofia University
5 J. Bourchier Blvd, 1164 Sofia, Bulgaria

August 6, 2012

Abstract

In this paper we consider the finite size effects for the strings in β -deformed $AdS_5 \times T^{1,1}$ background. We analyze the finite size corrections for the cases of giant magnon and single spike string solution. The finite size corrections for the undeformed case are straightforwardly obtained sending the deformation parameter to zero.

1 Introduction

After the remarkable AdS/CFT conjecture of Maldacena [1] relating the string theory on $AdS_5 \times S^5$ to the $\mathcal{N} = 4$ super Yang-Mills the search for less supersymmetric cases has started. One possible way is to consider a stack of N D3-branes at the tip of the conifold [2] resulting in the spacetime being the product $AdS_5 \times T^{1,1}$. The dispersion relations on the string theory side plays an important role, since they correspond to the large N limit of the anomalous dimension of particular gauge theory operators. Finding the spectrum is however a complicated task and one usually considers first states for which one of the string charges is infinite. This simplification can be seen in the light-cone gauge as a decompactifying limit, since one can rescale the worldsheet coordinate, such that the theory is defined on a cylinder with circumference proportional to the light-cone momentum (see [3] and references therein). In [4] it was shown that at large λ in the decompactifying limit a one-magnon excitation with the finite worldsheet momentum can be identified with the classical string sigma model. This string solution will have infinite momentum and infinite energy, but the difference of the two will be finite and equal to the energy of the worldsheet soliton [4, 3]. On the target space it corresponds to an open rigidly moving string with the distance of the endpoints fixed and proportional to the worldsheet momentum. This solution constructed in the conformal gauge was called *the giant magnon* [4]. The situation on the conifold was analyzed in [5].

One possible way of breaking the supersymmetry was found by Lunin and Maldacena in [6]. They considered a marginally deformed $\mathcal{N} = 4$ super Yang-Mills and found its gravity dual. This idea was further extended in [7, 8] and applied to other backgrounds, see [9].

Since all these considerations solve the theory on a plane, one can ask what is the difference to the physical case of a cylinder worldsheet. Such a modification can be significant, but for

^{*}e-mail: kanister@hep.itp.tuwien.ac.at.

^{\dagger}e-mail: rash@hep.itp.tuwien.ac.at.

sufficiently large size of the cylinder one can use an asymptotic construction [3] and obtain the *finite size corrections*.

In this paper we extend our previous considerations [9] computing the finite size corrections for the case of giant magnons and single spikes on the γ -deformed $T^{1,1}$. Since the deformation is continuous we obtain the finite size correction to the undeformed case [5] as well, simply by setting $\tilde{\gamma} = 0$.

2 A review of the case of infinite charges and finite size setup

In this section we review the results of the case of infinite charges [9], reformulate some of the important steps of the calculations to a convenient form and make a setup for the finite size calculations. Setting $\tilde{\gamma} = 0$ one can recover the undeformed case $AdS_5 \times T^{1,1}$ and the obtained results translate to that case.

2.1 Review of the infinite case

The TsT transformations [6, 8, 10] of the standard conifold metric results in the metric and the B-field of the γ -deformed $T^{1,1}$

$$\frac{ds^2}{R^2} = ds_{AdS}^2 + G \left(\frac{1}{6} \sum_{i=1}^2 (G^{-1} d\theta_i^2 + \sin^2 \theta_i d\phi_i^2) + \frac{1}{9} (d\psi + \cos \theta_1 d\phi_1 + \cos \theta_2 d\phi_2)^2 + \tilde{\gamma} \frac{\sin^2 \theta_1 \sin^2 \theta_2}{324} d\psi^2 \right), \quad (1)$$

$$\frac{B}{R^2} = \tilde{\gamma} G \left(\left(\frac{\sin^2 \theta_1 \sin^2 \theta_2}{36} + \frac{\cos^2 \theta_1 \sin^2 \theta_2 + \cos^2 \theta_2 \sin^2 \theta_1}{54} \right) d\phi_1 \wedge d\phi_2 + \frac{\sin^2 \theta_1 \cos \theta_2}{54} d\phi_1 \wedge d\psi - \frac{\cos \theta_1 \sin^2 \theta_2}{54} d\phi_2 \wedge d\psi \right). \quad (2)$$

The standard conifold metric is the one in (1) with $\tilde{\gamma} = 0$ and the conformal factor G in (1),(2) is

$$G^{-1} = 1 + \tilde{\gamma} \left(\frac{\cos^2 \theta_1 \sin^2 \theta_2 + \cos^2 \theta_2 \sin^2 \theta_1}{54} + \frac{\sin^2 \theta_1 \sin^2 \theta_2}{36} \right).$$

Since the complete $T^{1,1}$ background amounts to a complicated set of equations of motion we proceed with a consistent truncation to the subspace defined by

$$\theta_2 = 0, \quad \phi_2 = \text{const.}$$

The ansatz leading to a solitonic type solution is

$$t = \kappa \tau, \quad \theta_2 = 0, \quad \phi_2 = \text{const.} \\ \Psi = \omega_\psi \tau - \psi(y), \quad \Phi = \omega_\phi \tau + \phi(y), \quad \theta = \theta(y),$$

where $y = c\sigma - d\tau$ and setting $R = 1$ we restrict the metric (1) and the B-field (2) to

$$ds^2 = -dt^2 + \frac{1}{6} d\theta^2 + \frac{G}{6} \sin^2 \theta d\phi^2 + \frac{G}{9} (d\psi + \cos \theta d\phi)^2 \\ B = G \tilde{\gamma} \frac{\sin^2 \theta}{54} \\ G^{-1} = 1 + \tilde{\gamma} \frac{\sin^2 \theta}{54}. \quad (3)$$

The Lagrangian in the coordinates (3) is of the form

$$L = \dot{t}^2 + \frac{c^2 - d^2}{6} \theta'^2 + \frac{G}{9} \left(1 + \frac{\sin^2 \theta}{2} \right) \left(-(\omega_\phi - d\phi')^2 + c^2 \phi'^2 \right) + \\ \frac{2}{9} G \cos \theta \left(-(\omega_\psi - d\psi') (\omega_\phi - d\phi') + c^2 \psi' \phi' \right) + \frac{G}{9} \left(-(\omega_\psi - d\psi')^2 + c^2 \psi'^2 \right) + \\ 2G\tilde{\gamma} \frac{\sin^2 \theta}{54} c \left((\omega_\phi - d\phi') \psi' - (\omega_\psi - d\psi') \phi' \right)$$

and the once integrated equations of motion are (A_ϕ and A_ψ below are the integration constants)

$$(c^2 - d^2) \phi' + d\omega_\phi = \frac{3(A_\phi - A_\psi \cos \theta)}{G \sin^2 \theta} + \tilde{\gamma} \frac{c}{9} (\omega_\psi + \omega_\phi \cos \theta) \\ (c^2 - d^2) \psi' + d\omega_\psi = \frac{3(A_\psi - A_\phi \cos \theta)}{G \sin^2 \theta} + \frac{3A_\psi}{2G} - \frac{\tilde{\gamma}}{9} c \left(\omega_\psi \cos \theta + \omega_\phi \left(1 + \frac{\sin^2 \theta}{2} \right) \right).$$

The Virasoro constraints [9] read off

$$\frac{c^2 + d^2}{6} \theta'^2 + \frac{G}{9} \left(1 + \frac{\sin^2 \theta}{2} \right) \left((\omega_\phi - d\phi')^2 + c^2 \phi'^2 \right) + 2\frac{G}{9} \cos \theta \left((\omega_\psi - d\psi') (\omega_\phi - d\phi') + \right. \\ \left. c^2 \psi' \phi' \right) + \frac{G}{9} \left((\omega_\psi - d\psi')^2 + c^2 \psi'^2 \right) = \kappa^2,$$

$$\frac{1}{6} \theta'^2 + \frac{G}{9} \left(1 + \frac{\sin^2 \theta}{2} \right) \left(\phi'^2 - \frac{\omega_\phi \phi'}{d} \right) + \frac{G}{9} \left(\psi'^2 - \frac{\omega_\psi \psi'}{d} \right) + \\ \frac{G}{9} \cos \theta \left(2\phi' \psi' - \frac{\omega_\psi \phi' + \omega_\phi \psi'}{d} \right) = 0$$

and the restriction on the parameters is just

$$2d\kappa^2 = A_\phi \omega_\phi + A_\psi \omega_\psi. \quad (4)$$

Using the second Virasoro constraint we obtain the equation for θ

$$\theta'^2 + \frac{1}{3(c^2 - d^2)^2 \sin^2 \theta} \left(\frac{2c\omega_\phi - \tilde{\gamma}A_\psi}{2} \right)^2 \cos^4 \theta - (2c\omega_\phi - \tilde{\gamma}A_\psi) (2c\omega_\psi - \tilde{\gamma}A_\phi) \cos^3 \theta - \\ \frac{1}{2} \left((2c\omega_\psi + \tilde{\gamma}A_\phi)^2 + 2(2c\omega_\phi - \tilde{\gamma}A_\psi)^2 + 27A_\psi^2 - \frac{18}{d} (c^2 + d^2) (A_\phi \omega_\phi + A_\psi \omega_\psi) \right) \cos^2 \theta + \\ ((2c\omega_\phi - \tilde{\gamma}A_\psi) (2c\omega_\psi + \tilde{\gamma}A_\phi) - 54A_\phi A_\psi) \cos \theta + \frac{1}{2} \left((2c\omega_\psi + \tilde{\gamma}A_\phi)^2 + \frac{3}{2} (2c\omega_\phi - \tilde{\gamma}A_\psi)^2 + \right. \\ \left. 27(3A_\psi^2 + 2A_\phi^2) - \frac{18}{d} (c^2 + d^2) (A_\phi \omega_\phi + A_\psi \omega_\psi) \right) = 0. \quad (5)$$

2.2 Finite size analysis setup

In [9] we considered the case of infinite charges. In contrast to this case the finite size corrections originate from the expression

$$\int_0^{w_{max}} d\sigma = \int_0^{w_{max}} \frac{dy}{c} = \frac{1}{c} \int_{u(0)}^{u(w_{max})} \frac{du}{u'} \quad (6)$$

where $u = \cos^2 \left(\frac{\theta(y)}{2} \right)$ and the worldsheet coordinate σ is in the range $\langle -w_{max}, w_{max} \rangle$ with w_{max} finite. Rewriting the equation of motion (5) in $u(y)$ we obtain an equation accounting to the finite size

$$u'^2 = -f^2 (u - u_0) (u - u_1) (u - u_2) (u - u_3) = P_4(u). \quad (7)$$

The quartic polynomial $P_4(u)$ on the right hand side of (7) has in general four different roots u_i . In order to have a well defined derivative u' in the worldsheet integral (6) the integration has to be taken in the positive region of the $P_4(u)$. We are looking for certain string profiles, namely magnons and spikes, which have one turning point. This condition in the u coordinate is simply $u(w_{max}) = u_{turn} = u_j$ i.e. the turning point is one of the roots u_i . Our aim is to find finite size corrections to the dispersion relations obtained in the infinite case, therefore the limit $w_{max} \rightarrow \infty$ amounts to setting the turning point to $u_{turn} = 0$. Looking at the following relation

$$y = \int_{u(0)}^u \frac{d\tilde{u}}{u'} = F[u] - F[u(0)]$$

and the form of u' (7) it is clear that demanding $y \rightarrow \infty$ for $u_{turn} \rightarrow 0$ we actually demand $F[u] \xrightarrow{u_{turn} \rightarrow 0} \infty$. The latter is possible if there is another root u_k that equals u_{turn} at $w_{max} = \infty$. In order to have finite y for any $u \neq 0$ we demand one of the roots, say u_0 to be $u_0 = 0$. The polynomial $P_4(u)$ then changes to

$$P_4(u) = -f^2 u (u - u_1) (u - u_2) (u - u_3),$$

which results in the condition $A_\phi = -A_\psi$. Using the notations of [9]

$$B_\phi = 2c\omega_\phi + \tilde{\gamma}A_\phi, \quad B_\psi = 2c\omega_\psi + \tilde{\gamma}A_\phi \quad (8)$$

we find $P_4(u)$ in the following form

$$\begin{aligned} P_4(u) = & -\frac{1}{3(c^2 - d^2)^2} (B_\phi^2 u^4 - 2B_\phi(B_\phi + B_\psi)u^3 + \\ & \frac{1}{2} \left(B_\phi^2 - B_\psi^2 + 6B_\phi B_\psi - 27A_\phi^2 + \frac{18}{d} (c^2 + d^2) A_\phi (\omega_\phi - \omega_\psi) \right) u^2 + \\ & \frac{1}{2} \left((B_\psi - B_\phi)^2 + 81A_\phi^2 - \frac{18}{d} (c^2 + d^2) A_\phi (\omega_\phi - \omega_\psi) \right) u. \end{aligned} \quad (9)$$

The prefactor f^2 is thus

$$f^2 = \frac{B_\phi^2}{3(c^2 - d^2)^2}.$$

The solution to the equation (7) written in the implicit form is

$$\int_{u_3}^u \frac{d\tilde{u}}{\sqrt{|\tilde{u}(\tilde{u} - u_1)(\tilde{u} - u_2)(u_3 - \tilde{u})|}} = y,$$

where we set u_2 to be the turning point and $u_3 > u_2 > 0$. Analyzing the polynomial $P_4(u)$ one finds that $u_1 < 0$. In order to work with positive parameters u_i , it is convenient to define $u_{11} = -u_1$.

3 Finite size corrections

In this section we proceed with the conserved charges. Referring to [9] for further details, we start with the relations for conserved charges in the finite case

$$J_i = 2 \int_0^{w_{max}} \frac{dy}{c} P_i = \frac{2}{c} \int_{u_3}^{u_2} \frac{du}{u'} P_i.$$

Using the notations

$$I_1 = \int_{u_3}^{u_2} \frac{du}{u'}; I_2 = \int_{u_3}^{u_2} u \frac{du}{u'}; I_3 = \int_{u_3}^{u_2} u^2 \frac{du}{u'}; I_4 = \int_{u_3}^{u_2} \frac{1}{1-u} \frac{du}{u'} \quad (10)$$

we write the general expression for conserved charges

$$\begin{aligned} cE &= 2T\kappa I_1 \\ 9c(c^2 - d^2) J_\psi &= T(9dA_\phi - c(B_\phi - B_\psi)) I_1 + 2cTB_\phi I_2 \\ 9c(c^2 - d^2) J_\phi &= T(c(B_\phi - B_\psi) - 9dA_\phi) I_1 + 2cT(B_\phi + B_\psi) I_2 - 2cTB_\phi I_3 \\ 9c(c^2 - d^2) \Delta &= \frac{9}{2}(3cA_\phi - d(B_\phi - B_\psi)) I_1 + \tilde{\gamma}c((2B_\phi + B_\psi) I_2 - B_\phi I_3) + 27cA_\phi I_4. \end{aligned} \quad (11)$$

The integrals I_i are expressed in terms of complete elliptic integrals (34) and thus the above set of equations (11) becomes transcendental one. From the Virasoro constraints (4) we have

$$\kappa = \sqrt{\frac{A_\phi(\omega_\phi - \omega_\psi)}{2d}},$$

which results in

$$I_1 = \frac{c\sqrt{d}E}{T\sqrt{2A_\phi(\omega_\phi - \omega_\psi)}}. \quad (12)$$

In order to proceed we assume that our turning point u_2 is very small i.e. $u_2 \mapsto \epsilon \rightarrow 0$ which is equivalent to having large w_{max} and so large, but finite charges. In the rest of this paper we expand our calculations in $\epsilon \equiv u_2$. The turning point condition $u'(\epsilon) = 0 + O(\epsilon)^2$ gives again two relations for the parameter A_ϕ defining the magnon and the spike solution

$$A_\phi \begin{cases} \frac{2}{9}d(\omega_\phi - \omega_\psi) + \frac{1}{243(c^2 - d^2)(\omega_\phi - \omega_\psi)} \\ 4d((27c^2 + 12dc\tilde{\gamma} + d^2(\tilde{\gamma}^2 + 9))\omega_\phi^2 + d(d(\tilde{\gamma}^2 + 9) - 6c\tilde{\gamma})\omega_\psi^2 + & \text{magnon} \\ 2(27c^2 - 3dc\tilde{\gamma} - d^2(\tilde{\gamma}^2 + 9))\omega_\psi\omega_\phi) \epsilon + O(\epsilon^2) \\ \frac{2c^2}{9d}(\omega_\phi - \omega_\psi) + \frac{1}{243d(c^2 - d^2)(\omega_\phi - \omega_\psi)} \\ 4c^2((27d^2 + 12dc\tilde{\gamma} + c^2(\tilde{\gamma}^2 + 9))\omega_\phi^2 + c(c(\tilde{\gamma}^2 + 9) - 6d\tilde{\gamma})\omega_\psi^2 + & \text{slope} \\ 2(27d^2 - 3dc\tilde{\gamma} - d^2(\tilde{\gamma}^2 + 9))\omega_\psi\omega_\phi) \epsilon + O(\epsilon^2) \end{cases} \quad (13)$$

Since A_ϕ receives ϵ -correction so does B_ϕ, B_ψ , therefore for convenience we write them as

$$B_{\phi,\psi} = B_0 + \epsilon B_1 + O(\epsilon)^2. \quad (14)$$

Using the above form of parameters $B_{\phi,\psi}$ (14) and the expansions of the elliptic integrals (35) we

find the following expressions for the integrals I_i :

$$\begin{aligned}
I_1 &= \frac{\sqrt{3}(c^2 - d^2)}{B0_\phi \sqrt{u_{11}u_3}} \left(\log \left(\frac{16u_{11}u_3}{(u_{11} + u_3)\epsilon} \right) + \frac{1}{4u_{11}u_3} \right. \\
&\quad \left. \left(\left(u_{11} - u_3 - 4\frac{B1_\phi}{B0_\phi}u_{11}u_3 \right) \log \left(\frac{16u_{11}u_3}{(u_{11} + u_3)\epsilon} \right) - 2(u_{11} - u_3) \right) \epsilon + O(\epsilon)^2 \right) \\
I_2 &= \frac{\sqrt{3}(c^2 - d^2)}{B0_\phi} \left(2 \arctan \left(\sqrt{\frac{u_3}{u_{11}}} \right) + \right. \\
&\quad \left. \left(\frac{B0_\phi}{2\sqrt{3}(c^2 - d^2)} I_1 - \frac{1}{2\sqrt{u_{11}u_3}} - 2\frac{B1_\phi}{B0_\phi} \arctan \left(\sqrt{\frac{u_3}{u_{11}}} \right) \right) \epsilon \right) + O(\epsilon)^2 \\
I_3 &= \frac{\sqrt{3}(c^2 - d^2)}{B0_\phi} \left(\sqrt{u_{11}u_3} + (u_3 - u_{11}) \arctan \left(\sqrt{\frac{u_3}{u_{11}}} \right) + \right. \\
&\quad \left. \left(-\frac{B1_\phi}{B0_\phi} \sqrt{u_{11}u_3} + \left(1 + \frac{B1_\phi}{B0_\phi} (u_{11} - u_3) \right) \arctan \left(\sqrt{\frac{u_3}{u_{11}}} \right) \right) \epsilon \right) + O(\epsilon)^2 \\
I_4 &= I_1 + \frac{6(c^2 - d^2)}{\sqrt{3}B0_\phi \sqrt{(1 + u_{11})(1 - u_3)}} \arctan \left(\sqrt{\frac{(1 + u_{11})u_3}{u_{11}(1 - u_3)}} \right) + \left(\frac{1}{2} I_1 + \right. \\
&\quad \left. \frac{\sqrt{3}(c^2 - d^2)}{2B0_\phi \sqrt{u_{11}u_3}} \left(-1 + 2 \left(1 - 2\frac{B1_\phi}{B0_\phi} \right) \sqrt{\frac{u_{11}u_3}{(1 + u_{11})(1 - u_3)}} \right) \arctan \left(\sqrt{\frac{(1 + u_{11})u_3}{u_{11}(1 - u_3)}} \right) \right) \epsilon + O(\epsilon)^2
\end{aligned} \tag{15}$$

One see that the first integral I_1 is divergent for $\epsilon \rightarrow 0$ and behaves as $I_1|_{\epsilon \rightarrow 0} \sim -\log(\epsilon)$. Using this fact we get from the integral I_1 (15)

$$\frac{I_1 \sqrt{u_{11}u_3} B0_\phi}{\sqrt{3}(c^2 - d^2)} + \log \left(\frac{(u_{11} + u_3)\epsilon}{16u_{11}u_3} \right) + O(\epsilon^1) = 0. \tag{16}$$

Substituting I_1 from (12) and solving (16) we find

$$\epsilon \approx \frac{16e^{\mathcal{E}X} u_{11}u_3}{u_{11} + u_3}, \quad X = -\frac{B_\phi c \sqrt{d} \sqrt{u_{11}u_3}}{\sqrt{6}(c^2 - d^2) \sqrt{A_\phi(\omega_\phi - \omega_\psi)}}, \quad \mathcal{E} = E/T. \tag{17}$$

We can now return to the calculation of u_{11}, u_3 from (9). In order to expand in the ϵ parameter first we write

$$u_{11,3} = u_{11,3}^0 + \epsilon u_{11,3}^1. \tag{18}$$

The turning point condition simplifies $P_4(u) = 0$ (9) to the form

$$\begin{aligned}
&u(27A0_\phi^2 + (u - 1)B0_\phi(uB0_\phi - B0_\phi - 2B0_\psi)) + \epsilon(54uA0_\phi A1_\phi + 27(u - 1)A0_\phi^2 + (u - 1) \\
&(2u^2B0_\phi B1_\phi - 2u(B0_\psi B1_\phi + B0_\phi(B1_\psi + B1_\phi)) + B0_\phi(B0_\phi + 2B0_\psi)) + O(\epsilon^2) = 0,
\end{aligned} \tag{19}$$

where we expanded the parameter A_ϕ in ϵ as $A_\phi = A0_\phi + \epsilon A1_\phi$. The solution to (19) at the zeroth order in ϵ , which we call $u_{11,3}^0$, coincides with the infinite case results in [9]. Thus we have

$$u_3 = 1 + \frac{B0_\psi}{B0_\phi} + \frac{\sqrt{B0_\psi^2 - 27A0_\phi^2}}{B0_\phi} + \epsilon u_{13}^1, \quad -u_1 \equiv u_{11} = -1 - \frac{B0_\psi}{B0_\phi} + \frac{\sqrt{B0_\psi^2 - 27A0_\phi^2}}{B0_\phi} + \epsilon u_{11}^1 \tag{20}$$

with

$$\begin{aligned}
u_{13} &= \frac{-1}{2B0_\phi^2 \sqrt{B0_\psi^2 - 27A0_\phi^2}} (27A0_\phi^2 (B0_\phi - 2B1_\phi) + 54A0_\phi B0_\phi A1_\phi + \\
&\quad (2B0_\psi B1_\phi - 2B0_\phi B1_\psi + B0_\phi^2) (\sqrt{B0_\psi^2 - 27A0_\phi^2} + B0_\psi)) \\
u_{111} &= \frac{-1}{2B0_\phi^2 \sqrt{B0_\psi^2 - 27A0_\phi^2}} (27A0_\phi^2 (B0_\phi - 2B1_\phi) + 54A0_\phi B0_\phi A1_\phi - \\
&\quad (2B0_\psi B1_\phi - 2B0_\phi B1_\psi + B0_\phi^2) (\sqrt{B0_\psi^2 - 27A0_\phi^2} - B0_\psi))
\end{aligned} \tag{21}$$

In the further analysis we consider the different cases of giant magnon and single spike.

3.1 Giant magnon

We start with the value of A_ϕ (13) specifying the magnon solution. In order to find the dispersion relation we will substitute the integrals I_i from (15),(12) to the charges (11), use (18) as well as the relations

$$\begin{aligned}
Bm0_\psi &= -\frac{1}{2} (u_{011} - u_{03} + 2) Bm0_\phi \\
Bm1_\psi &= \frac{1}{2} ((-u_{011} + u_{03} - 2) Bm1_\phi + (-u_{111} + u_{13} + 1) Bm0_\phi) \\
\frac{Bm1_\phi}{Bm0_\phi} &= \frac{d\gamma (d^2 (u_{011} - u_{03} + 4)^2 - 12c^2 (u_{011} - u_{03} + 1))}{27c (c^2 - d^2) (u_{011} - u_{03} + 4)} + O(\epsilon).
\end{aligned}$$

The parameters BmI_α above are just $B0_{\phi,\psi}$, $B1_{\phi,\psi}$ from (14) with the value of A_ϕ defining the magnon case, i.e. $BmI_\alpha = BI_\alpha|_{A_\phi=A_{\phi\text{magnon}}}$, $I \in \{0, 1\}$, $\alpha \in \{\phi, \psi\}$. Thus we obtain relations for J_ϕ , J_ψ accounting for finite size effects

$$\begin{aligned}
\frac{4}{\sqrt{3}} \arctan \left(\sqrt{\frac{u_{03}}{u_{011}}} \right) &= \mathcal{E} + 3\mathcal{J}_\psi + \epsilon \left(\frac{2(u_{111}u_{03} - u_{13}u_{011})}{\sqrt{3}\sqrt{u_{011}u_{03}}(u_{011} + u_{03})} + \frac{1}{\sqrt{3}\sqrt{u_{011}u_{03}}} + \right. \\
&\quad \left. \frac{d^2 \mathcal{E} (d^2 (u_{011}^2 + (2 - 14u_{03})u_{011} + u_{03}^2 - 2u_{03} - 8) - 6c^2 (u_{011} - u_{03} - 2))}{36(c^2 - d^2)^2 (u_{011} + 1)(u_{03} - 1)} \right) + O(\epsilon^2) \tag{22}
\end{aligned}$$

and

$$\begin{aligned}
\frac{2\sqrt{u_{011}u_{03}}}{\sqrt{3}} &= (\mathcal{E} - 3\mathcal{J}_\phi) + \frac{\epsilon}{36((c^2 - d^2)^2 u_{011}(u_{011} + 1)(u_{03} - 1)u_{03})} (\\
&\quad \mathcal{E}u_{011}u_{03}d^2 (3c^2 ((u_{011} - u_{03})^2 - 4) - 2d^2 (u_{011}^2 + (4u_{03} + 2)u_{011} + (u_{03} - 4)(u_{03} + 2))) - \\
&\quad 6\sqrt{3}(c^2 - d^2)^2 (u_{011} + 1)(u_{03} - 1)\sqrt{u_{011}u_{03}}(2u_{111}u_{03} + (2u_{13} - 1)u_{011} + u_{03})) + O(\epsilon^2).
\end{aligned} \tag{23}$$

Above we used the charge densities $\mathcal{J}_i = J_i/T$ instead of charges. From the relations (22),(23) we find at the zeroth order approximation in ϵ

$$u_{011} = \frac{\sqrt{3}}{2} (\mathcal{E} - 3\mathcal{J}_\phi) \cot \left(\frac{\sqrt{3}}{4} (\mathcal{E} + 3\mathcal{J}_\psi) \right), \quad u_{03} = \frac{\sqrt{3}}{2} (\mathcal{E} - 3\mathcal{J}_\phi) \tan \left(\frac{\sqrt{3}}{4} (\mathcal{E} + 3\mathcal{J}_\psi) \right). \tag{24}$$

Substituting $u_{011,3}$ (24) to the angle difference equation we recover the infinite case dispersion relation at the zeroth order. At the first order we find the corrections $u_{111,3}$ which can be found in the appendix (37). The angle difference equation written in a concise form is

$$2 \arctan \left(\sqrt{\frac{(u_{011} + 1)u_{03}}{u_{011} - u_{011}u_{03}}} \right) = \Delta - \frac{1}{2} (\mathcal{J}_\phi + \mathcal{J}_\psi) \gamma + \frac{27\Delta_\epsilon}{c(c^2 - d^2)} \epsilon + O(\epsilon^2), \tag{25}$$

where $\Delta_\epsilon = \mathcal{E}\Delta_\mathcal{E} + \Delta_1$ and the parts $\Delta_\mathcal{E}, \Delta_1$ are given in the appendix (see 38). Since we assume $\epsilon \ll 1$ and thus the energy density $\mathcal{E}$ being very large, we see that the coefficient $\Delta_\mathcal{E}$ produces the leading part of the correction and the Δ_1 the subleading one. Expressing $\Delta - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma$ from (25) in a concise form [9] we obtain

$$\cos\left(\Delta - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma\right) = \frac{u0_{11} - u0_3 - 2u0_3u0_{11}}{u0_{11} + u0_3} + \frac{108\Delta_\epsilon u0_{11}u0_3(1 + u0_{11})(1 - u0_3)\epsilon}{c(c^2 - d^2)(u0_{11} + u0_3)^2 \sin\left(\Delta - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma\right)} + O(\epsilon^2). \quad (26)$$

Restricted to the first term it is just the infinite case dispersion relation.

In order to find the dispersion relation in terms of conserved charges we must substitute for the ϵ parameter from (17). Using the convenient notations $R_\phi = \frac{\sqrt{3}}{2}(\mathcal{E} - 3\mathcal{J}_\phi)$, $R_\psi = \frac{\sqrt{3}}{2}(\mathcal{E} + 3\mathcal{J}_\psi)$, $R_\Delta = \Delta - \frac{\gamma}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)$ we obtain the giant magnon dispersion relation (26) as

$$\cos(R_\Delta) = (\cos(R_\psi) - R_\phi \sin(R_\psi)) \left(1 + C_1 \mathcal{E} e^{\mathcal{E}X} + D_1 e^{\mathcal{E}X} + O(e^{\mathcal{E}X})^2\right). \quad (27)$$

The finite size correction consists of two parts $C_1 \mathcal{E} e^{\mathcal{E}X}$ and $D_1 e^{\mathcal{E}X}$ where C_1, D_1 are lengthy expressions of R_ϕ, R_ψ, R_Δ , that can be found in the appendix (39),(40). The exponent X in (27) is of the form

$$X = -\frac{2\sqrt{3}R_\phi \sin(R_\psi)(R_\phi \cos(2R_\psi) + 2\sin(R_\psi))}{1 + 4R_\phi^2 - (1 + 2R_\phi^2)\cos(2R_\psi) - 2R_\phi \sin(2R_\psi)} < 0,$$

or equivalently

$$X = -\frac{3c^2(\mathcal{E} - 3\mathcal{J}_\phi)}{(c^2 - d^2)\left(4 + \sqrt{3}(\mathcal{E} - 3\mathcal{J}_\phi)\cot\left(\frac{\sqrt{3}}{2}(\mathcal{E} + 3\mathcal{J}_\psi)\right)\right)},$$

which shows exponential suppression.

3.2 Single spike

For the case of a single spike defined by the value of A_ϕ (13) we use the same procedure as in the magnon case. In order to find the dispersion relation including the finite size correction we substitute the integrals I_i from (15),(12) to the relations for conserved charges (11) and use the relations :

$$\begin{aligned} Bs0_\psi &= -\frac{1}{2}(u0_{11} - u0_3 + 2)Bs0_\phi \\ Bs1_\psi &= \frac{1}{2}((-u0_{11} + u0_3 - 2)Bs1_\phi + (-u1_{11} + u1_3 + 1)Bs0_\phi) \\ \frac{Bs1_\phi}{Bs0_\phi} &= \frac{c\gamma(c^2(u0_{11} - u0_3 + 4)^2 - 12d^2(u0_{11} - u0_3 + 1))}{27d(c^2 - d^2)(u0_{11} - u0_3 + 4)} + O(\epsilon). \end{aligned}$$

Here $BsI_\alpha = BI_\alpha|_{A_\phi=A_{\phi spike}}$, $I \in \{0, 1\}$, $\alpha \in \{\phi, \psi\}$ from (14) and (13). Introducing the charge densities $\mathcal{J}_i = J_i/T$ we obtain the following expressions for the conserved charges J_ϕ, J_ψ (11) :

$$\begin{aligned} \frac{4}{\sqrt{3}} \arctan\left(\sqrt{\frac{u0_3}{u0_{11}}}\right) &= 3\mathcal{J}_\psi + \epsilon \left(\frac{c^3 \mathcal{E} ((u0_{11} - u0_3 + 1)(u0_{11} - u0_3 + 4)c^2 + 9d^2(u0_3 - u0_{11}))}{18d(c^2 - d^2)^2(u0_{11} + 1)(1 - u0_3)} \right. \\ &\quad \left. + \frac{\sqrt{u0_{11}u0_3}(-2u1_3u0_{11} + u0_{11} + 2u1_{11}u0_3 + u0_3)}{\sqrt{3}u0_{11}u0_3(u0_{11} + u0_3)} \right) + O(\epsilon^2) \quad (28) \end{aligned}$$

and

$$\mathcal{J}_\phi = -\frac{2\sqrt{u_{011}u_{03}}}{3\sqrt{3}} - \frac{1}{108}\epsilon \left(\frac{6\sqrt{3}(2u_{111}u_{03} + u_{03} + u_{011}(2u_{13} - 1))}{\sqrt{u_{011}u_{03}}} + \frac{c^3\mathcal{E}}{(c^2 - d^2)^2(u_{011} + 1)(1 - u_{03})} \left(2\sqrt{3}c(u_{011} - u_{03} - 8)\sqrt{(u_{011} + 1)(1 - u_{03})} - 3d(u_{011}^2 - 2(u_{03} + 2)u_{011} + u_{03}^2 + 4u_{03} - 8) \right) \right) + O(\epsilon^2) \quad (29)$$

Again, at the zeroth order in ϵ we find for the parameters $u_{011,3}$

$$u_{011} = -\frac{3}{2}\sqrt{3}\mathcal{J}_\phi \cot\left(\frac{3\sqrt{3}}{4}\mathcal{J}_\psi\right), \quad u_{03} = -\frac{3}{2}\sqrt{3}\mathcal{J}_\phi \tan\left(\frac{3\sqrt{3}}{4}\mathcal{J}_\psi\right) \quad (30)$$

which substituted to the remaining conserved quantity, the angle difference Δ , gives the infinite case dispersion relation. From the equations (28),(29) we calculate the corrections $u_{11,3}$, which are given in the appendix (see 41). Using (30),(41) we find

$$2 \arctan\left(\sqrt{\frac{(u_{011} + 1)u_{03}}{u_{011} - u_{011}u_{03}}}\right) = \Delta - \frac{3}{2}\mathcal{E} - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma + \frac{27\Delta_\epsilon}{c(c^2 - d^2)}\epsilon + O(\epsilon^2), \quad (31)$$

with $\Delta_\epsilon = \mathcal{E}\Delta_\mathcal{E} + \Delta_1$. The leading $\Delta_\mathcal{E}$ and the subleading Δ_1 part of Δ_ϵ are in the appendix (42). The other convenient form of (31) reads off

$$\cos\left(\Delta - \frac{3}{2}\mathcal{E} - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma\right) = \frac{u_{011} - u_{03} - 2u_{03}u_{011}}{u_{011} + u_{03}} + \frac{108\Delta_\epsilon u_{011}u_{03}(1 + u_{011})(1 - u_{03})\epsilon}{c(c^2 - d^2)(u_{011} + u_{03})^2 \sin\left(\Delta - \frac{3}{2}\mathcal{E} - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma\right)} + O(\epsilon^2), \quad (32)$$

with its zero-order being just the infinite case dispersion relation for single spike [9].

In order to find the explicit form of the corrections we substitute for ϵ from (17) and for $u_{011,3}$ from (30) and obtain

$$\cos\Delta\delta = \frac{3\sqrt{3}}{2}\left(\mathcal{J}_\phi \sin(R_\psi) + \frac{2\cos(R_\psi)}{3\sqrt{3}}\right)\left(1 + C_1\mathcal{E}e^{\mathcal{E}X} + D_1e^{\mathcal{E}X} + O(e^{\mathcal{E}X})^2\right). \quad (33)$$

Above we used $R_\psi = \frac{3\sqrt{3}}{2}\mathcal{J}_\psi$, $\Delta\delta = \Delta - \frac{3}{2}\mathcal{E} - \frac{1}{2}(\mathcal{J}_\phi + \mathcal{J}_\psi)\gamma$.

The expressions for C_1, D_1 can be found in the appendix (43),(44). The exponential factor X is of the form

$$X = -\sin^2(R_\psi) \frac{\mathcal{J}_\phi 9\sqrt{12 - 81\mathcal{J}_\phi^2 - 36\sqrt{3}\mathcal{J}_\phi \cot(R_\psi)}}{(2 + 27\mathcal{J}_\phi^2)\sin^2(R_\psi) + 3\mathcal{J}_\phi(9\mathcal{J}_\phi + 2\sqrt{3}\sin(2R_\psi))} < 0.$$

Since in this case $\mathcal{J}_\phi < 0$, negativity of the exponent is better recognizable from its equivalent form

$$X = \frac{6cd\mathcal{J}_\phi}{(c^2 - d^2)(3 - 2\sqrt{3}\mathcal{J}_\phi \cot(R_\psi))},$$

and we have again exponentially suppressed contribution.

4 Conclusion

In this paper we calculated the leading finite size corrections to the dispersion relations of giant magnon and single spike living on the γ -deformed Sasaki-Einstein manifold $T^{1,1}$. For the special case $\tilde{\gamma} = 0$ we get the finite size corrections to the undeformed case discussed in [5]. The result in this case has the same structure, with the difference in the R_Δ , $\Delta\delta$ parameters for the magnon and the spike case respectively. An interesting property of the results we obtained (27),(33) is the leading $\sim \mathcal{E}e^{\mathcal{E}X}$ and the subleading $\sim e^{\mathcal{E}X}$ part of the first order correction. Our results differs from the finite size correction for giant magnons in the conformal gauge on $\mathbb{R} \times S^2$ discussed in [3], which is natural from the reduced symmetry point of view. The authors of [3] considered finite size corrections of a one spin giant magnon, whereas we are discussing a two spin case. If we even restrict ourselves to the one spin case, which is a nontrivial limit of the results¹, we will obtain a correction for the consistent subsector of the $AdS_5 \times T^{1,1}$ which is quite different space than the maximally supersymmetric $AdS_5 \times S^5$. We restricted ourselves to the first order correction since higher orders behave extremely complicated. In principle however, one can carefully repeat the same steps and find corrections of higher order.

A Solution to the integrals and expansions

Using the following convenient notation $k = \frac{(u_0 - u_1)(u_3 - u_2)}{(u_1 - u_2)(u_0 - u_3)}$, $m = \frac{u_3 - u_2}{u_1 - u_2}$ we obtain the explicit solutions to the integrals I_i :

$$\begin{aligned}
I_1 &= -\frac{2\mathbb{K}(k)}{\sqrt{(u_1 - u_2)(u_0 - u_3)}} \\
I_2 &= -2\sqrt{\frac{1}{(u_1 - u_2)(u_0 - u_3)}}(u_1\mathbb{K}(k) + (u_3 - u_1)\mathbb{PI}(m, k)) \\
I_3 &= \frac{1}{\sqrt{(u_1 - u_2)(u_0 - u_3)}}\left(- (u_1(u_1 + u_2) + (u_1 - u_2)u_3)\mathbb{K}(k) - (u_1 - u_2)(u_0 - u_3)\mathbb{E}(k) \right. \\
&\quad \left. + (u_1 - u_3)(u_0 + u_1 + u_2 + u_3)\mathbb{PI}(m, k)\right) \\
I_4 &= -\frac{2\sqrt{\frac{u_1 - u_2}{u_0 - u_3}}\left((u_3 - 1)\mathbb{K}(k) + (u_1 - u_3)\mathbb{PI}\left(\frac{1 - u_1}{1 - u_3}m, k\right)\right)}{(u_1 - 1)(u_1 - u_2)(u_3 - 1)}
\end{aligned} \tag{34}$$

We use the following expansions for small ϵ :

$$\begin{aligned}
\mathbb{K}(1 - \epsilon) &= \left(2\log(2) - \frac{\log(\epsilon)}{2}\right) + \frac{1}{8}(-\log(\epsilon) + 4\log(2) - 2)\epsilon - \frac{21\epsilon^2}{128} - \frac{185\epsilon^3}{1536} + O(\epsilon^4) \\
\mathbb{E}(1 - \epsilon) &= 1 + \frac{1}{4}(-\log(\epsilon) + 2\log(4) - 1)\epsilon + \frac{1}{64}(-6\log(\epsilon) + 24\log(2) - 13)\epsilon^2 + \\
&\quad \frac{3}{256}(-5\log(\epsilon) + 20\log(2) - 12)\epsilon^3 + O(\epsilon^4) \\
\mathbb{PI}(m, 1 - \epsilon) &= \frac{2\sqrt{m}\operatorname{arctanh}(\sqrt{m}) + \log\left(\frac{\epsilon}{16}\right)}{2m - 2} + \frac{(-4\sqrt{m}\operatorname{arctanh}(\sqrt{m}) + (m + 1)\log\left(\frac{16}{\epsilon}\right) - 2)\epsilon}{8(m - 1)^2} \\
&\quad + \left((12 - 5m)m + 3(m - 6)\log\left(\frac{16}{\epsilon}\right)m + 48\operatorname{arctanh}(\sqrt{m})\sqrt{m} + \right. \\
&\quad \left. 9\log(\epsilon) - 36\log(2) + 21\right)\frac{\epsilon^2}{128(m - 1)^3} + O(\epsilon^4)
\end{aligned} \tag{35}$$

¹One must carefully take the limit $\omega_\psi \rightarrow 0$ and not just turn off the charge $\mathcal{J}_\psi$.

B Results and useful relations

B.1 The giant magnon case

Using the convenient notation $\left\{ X1 = \sqrt{u0_{11}u0_3}, X2 = \sqrt{\frac{u0_3}{u0_{11}}} \right\}$ we write the corrections $u1_{11,3}$ as

$$\begin{aligned}
(c^2 - d^2)^2 u1_{11} &= \frac{\mathcal{E}c^2d^2(-X1X2^2 + X1 - 2X2)}{4\sqrt{3}X2^2(X1X2 - 1)} - \frac{c^4}{2} + c^2d^2 - \frac{d^4}{2} - \\
&\frac{d^4\mathcal{E}(-16X1X2^4 + 2X1^2(X2^4 + 8X2^2 + 3)X2 + X1^3(X2^6 - 13X2^4 - 13X2^2 + 1) - 32X2^3)}{48\sqrt{3}X2^3(X1 + X2)(X1X2 - 1)} \\
(c^2 - d^2)^2 u1_3 &= \frac{c^2d^2(2X2(\mathcal{E}X2 - 2\sqrt{3}) + X1(\mathcal{E}X2(X2^2 - 1) - 4\sqrt{3}))}{4\sqrt{3}(X1 + X2)} + \frac{c^4}{2} + \frac{d^4}{2} + \\
&\frac{d^4\mathcal{E}(X1^3(X2^6 - 13X2^4 - 13X2^2 + 1) - 2X1^2(3X2^5 + 8X2^3 + X2) - 16X1X2^2 + 32X2^3)}{48\sqrt{3}X2(X1 + X2)(X1X2 - 1)}
\end{aligned} \tag{37}$$

The correction term $\Delta_\epsilon = \mathcal{E}\Delta_\mathcal{E} + \Delta_1$ in (25) consists of

$$\begin{aligned}
\Delta_\mathcal{E} &= \frac{d^3Z\sqrt{u0_{11}u0_3}}{324((u0_{11} + 1)(1 - u0_3))^{3/2}(u0_{11}^2 + 2(5u0_3 - 2)u0_{11} + (u0_3 + 2)^2) \\
&\quad (u0_{11}^4 + (8 - 12u0_3)u0_{11}^3 + (-26u0_3^2 + 8u0_3 - 36)u0_{11}^2 - \\
&\quad 4(3u0_3^3 + 2u0_3^2 + 26u0_3 - 4)u0_{11} + (u0_3 + 2)^2(u0_3^2 - 12u0_3 + 8))} \\
\Delta_1 &= \frac{1}{324(u0_{11} + 1)^2(u0_3 - 1)} \\
&\quad (2Z(u0_{11} + 1)(c^3(u0_{11}(9 - 6u0_3) - 9u0_3 + 12) - \\
&\quad cd^2(-11u0_3 + u0_{11}(2u0_3 + 11) + 20) + 4\sqrt{3}d^3\sqrt{-(u0_{11} + 1)(u0_3 - 1)}) + \\
&\quad \frac{\sqrt{3}d(c^2 - d^2)((u0_{11} - u0_3 + 5)(u0_{11}u0_3)^{3/2} + 4\sqrt{u0_{11}u0_3^3} - \sqrt{u0_{11}u0_3^5})}{u0_3}) \\
Z &= \arctan\left(\sqrt{\frac{(u0_{11} + 1)u0_3}{u0_{11} - u0_{11}u0_3}}\right).
\end{aligned} \tag{38}$$

The coefficients of the leading (C_1) and the subleading (D_1) part of the correction take the form:

$$\begin{aligned}
C_1 &= 8\sqrt{3}R_\Delta \csc(R_\Delta) R_\phi^2 \sin^3(R_\psi) (-2R_\phi \cos(R_\psi) - (1 - R_\phi^2) \sin(R_\psi)) (3 - 21R_\phi^2 - 4R_\phi^4 + \\
&\quad (-4 + 22R_\phi^2 + 8R_\phi^4) \cos(2R_\psi) + (1 - R_\phi^2) \cos(4R_\psi) + 8R_\phi \cos(R_\psi) \sin^3(R_\psi) + 8R_\phi^3 \sin(2R_\psi)) / \\
&\quad \left((-1 - 4R_\phi^2 + (1 + 2R_\phi^2) \cos(2R_\psi) + 2R_\phi \sin(2R_\psi))^2 \right. \\
&\quad \left. (R_\phi - 3R_\phi \cos(2R_\psi) + (-2 + R_\phi^2) \sin(2R_\psi)) \right)
\end{aligned} \tag{39}$$

$$\begin{aligned}
D_1 = & \frac{4R_\phi \csc(R_\Delta) \sin^2(R_\psi)}{-4Y(R_\phi \cos(R_\psi) + 2 \sin(R_\psi))(2R_\phi \sin(2R_\psi) + (2R_\phi^2 + 1) \cos(2R_\psi) - 4R_\phi^2 - 1)} \\
& (R_\phi \sin(R_\psi) - \cos(R_\psi))^{-1} (R_\phi^2 \sin(2R_\psi) (-22Y R_\Delta R_\phi - 8R_\phi^4 + 38R_\phi^2 + 2) + \\
& R_\phi^2 \sin(4R_\psi) (2R_\phi^4 - 2Y R_\Delta R_\phi^3 - 17R_\phi^2 + 7Y R_\Delta R_\phi - 1) + \\
& 4 \cos(2R_\psi) (R_\phi (R_\phi (R_\phi (-5Y R_\Delta R_\phi + 8R_\phi^2 - 4) + 2Y R_\Delta) - 2) - 2Y R_\Delta) + \\
& \cos(4R_\psi) (Y R_\Delta (8R_\phi^4 + R_\phi^2 + 2) + 2R_\phi (-5R_\phi^4 + 5R_\phi^2 + 1)) + \\
& 36Y R_\Delta R_\phi^4 - 9Y R_\Delta R_\phi^2 + 6Y R_\Delta - 6R_\phi^5 + 6R_\phi^3 + 6R_\phi)
\end{aligned} \tag{40}$$

with $Y = \sqrt{2R_\phi \cot(R_\psi) - R_\phi^2 + 1}$.

B.2 The single spike case

We find the the following form for $u_{11,3}$

$$\begin{aligned}
u_{11} = & -\frac{cd\mathcal{E}u_{011}}{2\sqrt{3}(c^2 - d^2)^2(u_{011} - u_{03} + 4)^2\sqrt{u_{011}u_{03}}} \left(c^2(-u_{03}^2 + u_{011}(u_{011} + 2) - 8) \right. \\
& \left. (u_{011} - u_{03} + 4) + 6d^2(-2u_{011}^2 + (u_{03} + 2)u_{011} + (u_{03} - 2)u_{03} + 4) \right) - \frac{1}{2} \\
u_{13} = & \frac{cd\mathcal{E}u_{03}}{2\sqrt{3}(c^2 - d^2)^2(u_{011} - u_{03} + 4)^2\sqrt{u_{011}u_{03}}} \left(c^2(u_{011}^2 - u_{03}^2 + 2u_{03} + 8) \right. \\
& \left. (u_{011} - u_{03} + 4) - 6d^2(u_{011} - u_{03})(u_{011} + 2u_{03} + 2) - 24d^2 \right) + \frac{1}{2}
\end{aligned} \tag{41}$$

The parameters $\Delta_{\mathcal{E}}$, Δ_1 read of

$$\begin{aligned}
\Delta_{\mathcal{E}} = & \frac{-4d^3 Z \sqrt{u_{011}u_{03}}}{3\sqrt{3}(u_{011} - u_{03} + 4)^4(u_{011}^2 + 2(5u_{03} - 2)u_{011} + (u_{03} + 2)^2)} \\
& ((4u_{03} - 3)u_{011}^4 + (4u_{03}^2 + 4u_{03} + 2)u_{011}^3 + \\
& 2(-2u_{03}^3 + 7u_{03}^2 + u_{03} + 12)u_{011}^2 - 2(2u_{03}^4 - 2u_{03}^3 + u_{03}^2 - 40u_{03} + 12)u_{011} - \\
& (u_{03} + 2)^2(3u_{03}^2 - 10u_{03} + 4)) \\
\Delta_1 = & \frac{1}{27(u_{011} - u_{03} + 4)^2} \left(\sqrt{3}d(d^2 - c^2)\sqrt{u_{011}u_{03}}(u_{011} - u_{03} + 4) + \right. \\
& \frac{2Z}{c} (2c^4(u_{011} - u_{03} + 4) - c^2d^2(-11u_{03} + u_{011}(2u_{03} + 11) + 20) + \\
& \left. 3d^4(u_{011}(3 - 2u_{03}) - 3u_{03} + 4)) \right) \\
Z = & \arctan \left(\sqrt{\frac{(u_{011} + 1)u_{03}}{u_{011} - u_{011}u_{03}}} \right).
\end{aligned} \tag{42}$$

The correction coefficients C_1, D_1 for the single spike dispersion relation are of the form

$$\begin{aligned}
C_1 = & 162\mathcal{J}_\phi^2 \Delta \delta \csc(\Delta \delta) \sqrt{4 - 27\mathcal{J}_\phi^2 - 12\sqrt{3}\mathcal{J}_\phi \cot(R_\psi) \sin^4(R_\psi) (12 - 810\mathcal{J}_\phi^2 + 1458\mathcal{J}_\phi^4 + \\
& (-16 + 864\mathcal{J}_\phi^2 + 2916\mathcal{J}_\phi^4) \cos(2R_\psi) + \sqrt{3}(-36\mathcal{J}_\phi + 162\mathcal{J}_\phi^3 + 2187\mathcal{J}_\phi^5) \sin(2R_\psi) \\
& + (4 - 54\mathcal{J}_\phi^2) \cos(4R_\psi) + 18\sqrt{3}\mathcal{J}_\phi \sin(4R_\psi)} \Bigg/ \left(\left(3\sqrt{3}\mathcal{J}_\phi \cos(R_\psi) - 4 \sin(R_\psi) \right) \right. \\
& \left. \left(2 \cos(R_\psi) + 3\sqrt{3}\mathcal{J}_\phi \sin(R_\psi) \right) \left(2 + 54\mathcal{J}_\phi^2 - (2 + 27\mathcal{J}_\phi^2) \cos(2R_\psi) + 6\sqrt{3}\mathcal{J}_\phi \sin(2R_\psi) \right)^2 \right)
\end{aligned} \tag{43}$$

$$\begin{aligned}
D_1 = & \frac{-3\mathcal{J}_\phi \csc(\Delta\delta) \sin^2(R_\psi) (3\sqrt{3}\mathcal{J}_\phi \sin(R_\psi) + 2\cos(R_\psi))^{-1}}{4(3\sqrt{3}\mathcal{J}_\phi \cos(R_\psi) - 4\sin(R_\psi)) (6\sqrt{3}\mathcal{J}_\phi \sin(2R_\psi) - (27\mathcal{J}_\phi^2 + 2)\cos(2R_\psi) + 54\mathcal{J}_\phi^2 + 2)} \\
& \left(8 \left(\sqrt{3}\Delta\delta (-3645\mathcal{J}_\phi^4 + 216\mathcal{J}_\phi^2 - 32) + 9Y\mathcal{J}_\phi (81\mathcal{J}_\phi^2 + 4) \right) \cos(2R_\psi) + \right. \\
& 2 \left(4\sqrt{3}\Delta\delta (27(54\mathcal{J}_\phi^4 + \mathcal{J}_\phi^2) + 8) - 9Y\mathcal{J}_\phi (81\mathcal{J}_\phi^2 + 4) \right) \cos(4R_\psi) + 6 \left(27\mathcal{J}_\phi^2 \sin(2R_\psi) \right. \\
& \left. \left(\sqrt{3}Y ((2 - 9\mathcal{J}_\phi^2) \cos(2R_\psi) + 18\mathcal{J}_\phi^2 - 2) + 6\Delta\delta\mathcal{J}_\phi ((27\mathcal{J}_\phi^2 - 14) \cos(2R_\psi) + 22) \right) - \right. \\
& \left. \left. 9Y(81\mathcal{J}_\phi^2 + 4)\mathcal{J}_\phi + 4\sqrt{3}\Delta\delta(2187\mathcal{J}_\phi^4 - 81\mathcal{J}_\phi^2 + 8) \right) \right) \quad (44)
\end{aligned}$$

where we used the notation $Y = \sqrt{-12\sqrt{3}\mathcal{J}_\phi \cot(R_\psi) - 27\mathcal{J}_\phi^2 + 4}$.

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.*, vol. 2, pp. 231–252, 1998.
- [2] I. R. Klebanov and E. Witten, “Superconformal field theory on threebranes at a Calabi-Yau singularity,” *Nucl. Phys.*, vol. B536, pp. 199–218, 1998.
- [3] G. Arutyunov, S. Frolov, and M. Zamaklar, “Finite-size Effects from Giant Magnons,” *Nucl. Phys.*, vol. B778, pp. 1–35, 2007.
- [4] D. M. Hofman and J. M. Maldacena, “Giant Magnons,” *J. Phys. A*, vol. A39, pp. 13095–13118, 2006.
- [5] S. Benvenuti and E. Tonni, “Giant magnons and spiky strings on the conifold,” *JHEP*, vol. 02, p. 041, 2009.
- [6] O. Lunin and J. M. Maldacena, “Deforming field theories with U(1) x U(1) global symmetry and their gravity duals,” *JHEP*, vol. 0505, p. 033, 2005.
- [7] R. G. Leigh and M. J. Strassler, “Exactly marginal operators and duality in four-dimensional N=1 supersymmetric gauge theory,” *Nucl. Phys.*, vol. B447, pp. 95–136, 1995.
- [8] S. Frolov, “Lax pair for strings in Lunin-Maldacena background,” *JHEP*, vol. 0505, p. 069, 2005.
- [9] H. Dimov, M. Michalcik, and R. C. Rashkov, “Strings on the deformed $T^{1,1}$: giant magnon and single spike solutions,” *JHEP*, vol. 10, p. 019, 2009.
- [10] A. Catal-Ozer, “Lunin-Maldacena deformations with three parameters,” *JHEP*, vol. 0602, p. 026, 2006.